

SOP: M001

Preparation of GAS medium protocol

Materials and Reagents:

1. Milli-Q water
2. Beaker, 1 liter
3. Magnetic stir bar
4. Magnetic stir plate
5. BactoCasitone (BD Science BD225930)
6. Ferric ammonium citrate (Sigma F-5879)
7. Potassium phosphate, dibasic anhydrous (VWR MK709208) (note 1)
8. Citric acid, anhydrous (VWR JT0122-1)
9. L-Alanine (Sigma A-7627)
10. Magnesium chloride, heptahydrate (VWR MK595804)
11. Potassium sulfate (VWR MK714004)
12. Ammonium chloride (VWR MK338412)
13. Sodium hydroxide, 10 M
14. Glycerol (VWR IC800689)
15. Graduated cylinder, 1 liter
16. Autoclave

Protocol:

1. ____ Pour 800 ml of Milli-Q water into a 1 liter beaker.
2. ____ Add magnetic stir bar to beaker and place on magnetic stir plate.
3. ____ Add 0.3 g of BactoCasitone.
4. ____ Add 0.05 g of ferric ammonium citrate.
5. ____ Add 4.0 g of potassium phosphate, dibasic anhydrous (note 1).
6. ____ Add 2.0 g of citric acid.
7. ____ Add 1.0 g of L-alanine.
8. ____ Add 1.2 g of magnesium chloride.
9. ____ Add 0.6 g of potassium sulfate.
10. ____ Add 2.0 g of ammonium chloride.
11. ____ Make sure all components are completely in solution.
12. ____ Add 1.8 ml of 10 M sodium hydroxide.
13. ____ Make sure the sodium hydroxide is completely in solution.
14. ____ Add 10.0 ml of glycerol.
15. ____ Make sure the glycerol is fully dispersed.
16. ____ Measure the pH, and adjust to 6.6

17. _____ Pour medium into 1 liter graduated cylinder.
18. _____ Bring volume to 1 liter with Milli-Q water.
19. _____ Transfer/aliquot to desired container(s).
20. _____ Autoclave on liquid cycle (slow exhaust) at 121°C for 15 minutes.

Notes:

1. **MUST** use Mallinckrodt K_2HPO_4 ; other brands will salt out during sterilization.

Reference:

Takayama, K., H. K. Schnoes, E. L. Armstrong, and R. W. Boyle. 1975. Site of inhibitory action of isoniazid in the synthesis of mycolic acids in *Mycobacterium tuberculosis*. *J. Lipid Res.* 16: 308-317.